

“Fermi Liquid” Shell Model Approach to Composite Fermion Excitation Spectra in Fractional Quantum Hall States

P. Sitko^{1,2}, S. N. Yi^{1,3}, K. S. Yi^{1,4}, and J. J. Quinn^{1,5}

¹*University of Tennessee, Knoxville, Tennessee 37969, USA*

²*Institute of Physics, Technical University of Wrocław, Wyb. Wyspiańskiego 27, 50-370 Wrocław, Poland*

³*Korea Maritime University, Pusan 606-791, Korea*

⁴*Pusan National University, Pusan 609-735, Korea*

⁵*Oak Ridge National Laboratory, Oak Ridge, Tennessee, 37831, USA*

(February 1, 2008)

Numerical results for the energy spectra of N electrons on a spherical surface are used as input data to determine the quasiparticle energies and the pairwise “Fermi liquid” interactions of composite Fermion (CF) excitations in fractional quantum Hall systems. The quasiparticle energies and their interactions are then used to determine the energy spectra, E vs total angular momentum L , of states containing more than two quasiparticles. The qualitative agreement with the numerical results gives a remarkable new confirmation of the CF picture.

PACS numbers: 73.40.Hm, 73.20.Dx

Numerical diagonalization of the interaction Hamiltonian for N electrons on a spherical surface in the presence of a radial magnetic field has contributed greatly to our understanding of the elementary excitations in fractional quantum Hall systems [1]. The energy spectra (E vs L , the total angular momentum) always show a lowest energy sector (or lowest band) separated by a well-defined gap from a first excited sector [2]. In many cases the first excited sector is separated from higher states by a slightly less well-defined gap. The object of the present paper is to demonstrate the extent to which the low lying states can be understood in terms of CF excitations [3,4] and their interaction energies.

We do not attempt to evaluate microscopically the quasiparticle energy (i.e. the energy of a single CF excitation interacting with a Laughlin condensed state) or the energy of interaction of a quasiparticle pair. Instead we determine these phenomenologically by using the exact numerical results for N -electron systems containing zero, one or two quasiparticles. These phenomenological “Fermi-liquid” parameters are then used to study states containing three or more quasiparticles within a shell model analogous to that of the nucleus [5]. The qualitative agreement of this “Fermi liquid” shell model approach with the exact numerical spectra gives striking new confirmation of the CF picture of the fractional quantum Hall effect. The small quantitative discrepancies may give some clue to the relative importance of the three body forces that arise in the Chern-Simons transformation from electrons to composite Fermions [4,6].

The single electron states on a Haldane sphere have eigenfunctions $|l, m\rangle$ and eigenvalues [7]:

$$\xi_l = \frac{\hbar\omega_c}{2S}(l(l+1) - S^2), \quad (1)$$

where l is the angular momentum, m its z -component, and $2S\frac{\hbar c}{e}$ is the strength of the magnetic monopole which produces the radial magnetic field. The allowed values of

l begin with the integral or half-integral value S and increase in unit steps. If we concentrate on a partially filled lowest Landau level ($l = S$), we have $2S + 1$ degenerate single particle states with m -values going from $-S$ to S . We can form $(2S + 1)[N!(2S + 1 - N)!]^{-1}$ antisymmetric many body states of N electrons from these degenerate single particle states. The many body states can be written $|\alpha, L, M\rangle$, where L and M are the total angular momentum and its z -component, and α distinguishes different states of the same L . The Wigner-Eckart theorem simplifies the diagonalization of the interaction Hamiltonian enormously, and the final energies depend upon L and α but not upon M .

In making the transformation from electrons to composite Fermions [3], an even number of fictitious flux quanta are attached to each electron. In the mean-field approximation, this leads to an effective magnetic field B^* and an effective CF filling factor ν^* which is an integer. Since we will concentrate on states close to the Laughlin $\nu = 1/3$ state ($\nu^* < 2$), we will make use of the result that for such states $2S = 3(N - 1) + n_{QH} - n_{QE}$ and $2S^* = 1(N - 1) + n_{QH} - n_{QE}$. Here n_{QH} (n_{QE}) is the number of CF quasihole (quasielectron) excitations. S^* plays exactly the same role for CF's that S plays for electrons [8], so that the angular momentum of a quasihole is $l_{QH} = S^*$ and of a quasielectron (which occurs in the first excited Landau level) $l_{QE} = S^* + 1$.

The low-lying states can be classified by the number of quasiparticle excitations they contain. The lowest energy sector contains the minimum number consistent with the values of N and $2S$. The first excited sector contains one additional QE-QH pair. The angular momentum of a state is simply the total angular momentum of the quasiparticle excitations [8]. This can be obtained by addition of angular momentum of the QE excitations and the QH excitations treated as independent sets of Fermions. The energy ξ_{QE} or ξ_{QH} of a single quasihole or a sin-

gle quasielectron can be obtained by subtracting the energy $E_{L=0}(2S_{\nu=1/3})$ from $E_{L=N/2}(2S = 2S_{\nu=1/3} \pm 1)$. A state containing n_{QE} quasielectrons and n_{QH} quasiholes would have energy $n_{QE}\xi_{QE} + n_{QH}\xi_{QH}$ if there were no interaction among the quasiparticle excitations. By looking at the numerical results it is apparent that the lowest energy sector of the $n_{QE} = 2$ or $n_{QH} = 2$ states and the first excited sector of the Laughlin condensed state contain a number of states which are not degenerate. The difference between the numerical results for the two quasiparticle state $|\alpha, L, M\rangle$ and the sum of the bare quasiparticle energies can be interpreted as the Fermi liquid interaction energy of the quasiparticles in that state [9]. For a QE-QH pair a plot of $V_{QE-QH}(L)$ for $N = 5, 6, 7, 8$ electron systems is given in reference [9]. For a QE-QE (or a QH-QH) pair it is more useful to plot V_{QE-QE} (or V_{QH-QH}) as a function of $L_{MAX} - L$, where L_{MAX} is the largest possible angular momentum of the pair [9,10]. The functions $V_{QP-QP'}(L)$ play the role of Landau's Fermi liquid interaction energies [11]. It should be noted that both ξ_{QP} and $V_{QP-QP'}(L)$ must contain effects due to both two particle and three particle interactions. ξ_{QP} is defined as the sum of the QP kinetic energy and its interaction with the Laughlin condensed state, which plays the role of the vacuum state. In the CF picture this interaction contains both two body and three body interaction terms.

Let us consider the system of 3 particles, two of them with angular momentum l , the third with angular momentum l' . The wave function of the system in the state of the total angular momentum L can be written using the Racah decomposition [12]:

$$|l^2(L_1)l'L\rangle = \sum_{L'} R_{L',L_1} |l'(L')lL\rangle. \quad (2)$$

Here $|l^2(L_1)l'L\rangle$ denotes the state in which two QP's of angular momentum l are coupled to give a resulting two particle angular momentum L_1 ; L_1 is then coupled with l' to give a total angular momentum L . In Eq. (2)

$$R_{L',L_1} = \sqrt{(2L_1+1)(2L'+1)} W(lLl'; L_1L'),$$

where W are Racah coefficients. The matrix element of the interaction energy is

$$\langle l^2(L_1)l'L | \sum_{i<k} V_{ik} | l^2(L_2)l'L \rangle.$$

We separate the interaction in the same shell (ll) and the interactions between shells ll' [5]; then this matrix element becomes

$$V_{ll}(L_1)\delta_{L_1,L_2} + 2 \sum_{L'} R_{L',L_1}^* R_{L',L_2} V_{ll'}(L'). \quad (3)$$

In the case of degenerate states the interaction energies are eigenvalues of the interaction matrix.

The problem of three or more identical particles all with angular momentum l involves the idea of coefficients of fractional parentage. In the special case of three particles we may write [12]:

$$|l^3L\alpha\rangle = \sum_{L'} F_{L'}^\alpha |l^2(L')lL\rangle$$

where $|l^2(L')lL\rangle$ is the wave function of the system in which the pair angular momentum $l(L')$ is defined and α is the additional parameter distinguishing different degenerate states. $F_{L'}^\alpha$ are coefficients of fractional parentage which satisfy the normalization condition $\sum_{L'} |F_{L'}^\alpha|^2 = 1$. The antisymmetry of the states leads to the matrix equation [12]: $\sum_{L''} F_{L''} (A_{L'',L'} - \delta_{L'',L'}) = 0$, where $A_{L'',L'} = (-)^{3l-L+1} \sqrt{(2L''+1)(2L'+1)} W(lLlL; L''L')$.

The matrix element of the interaction of the system is given by [5]:

$$\langle l^3L\alpha | \sum_{i<k} V_{ik} | l^3L\beta \rangle = 3 \sum_{L'} F_{L'}^\alpha F_{L'}^\beta V_{ll}(L'), \quad (4)$$

where $V_{ll}(L')$ is the pair interaction of particles in $l-l$ coupling in the state of the pair angular momentum L' .

The equations (3) and (4) allow us to obtain the energy of three particle states in terms of two-particle interactions. In our approach we make use of the quasiparticle picture of the shell theory of the nucleus [5]. The main idea is that the shell with $2l+1-n$ particles can be treated in terms of the conjugated configuration of l^n quasiholes. The allowed angular momenta of these two configurations are the same. The idea of quasiholes can be extended to the configuration of particles in two shells. For example the $l^{2l'}$ configuration can be considered as a particle-hole pair. One can treat the particle-hole interaction $V_{2l'l'}$ as the interaction between two particles ll' , $V_{ll'}$ [5].

Let us start by analyzing the spectrum of a three quasielectron excitation. We keep the number of electrons fixed and change the flux going through the sphere ($2S$); a decrease of one flux quantum from the Laughlin state generates one quasielectron of the angular momentum $l_{QE} = N/2$. A decrease of an additional flux quantum creates two quasielectrons in the angular momentum shell $l = N/2 - 1/2$. Hence, for $N=7$ the allowed angular momenta of a quasielectron pair are 5, 3, 1. In the next step we find the three-quasielectron excitations corresponding to three CF's in the shell $l = 2.5$. The construction of antisymmetric states leads to the states of the total angular momenta 1.5, 2.5, 4.5. In this configuration the pair angular momenta are 4, 2, 0, so that to find interaction energies of the system we have to know pairwise interactions for the shell $l = 2.5$. In fact, those can be found in the system with a different number of electrons, in this case for $N = 6$. Another possibility is to use the data for pairwise interaction for $N = 7$ assuming

that quasielectron-quasielectron interaction is a function of $RAM = L_{MAX} - L_{pair}$ [9,10]. Here we use the latter approach since the interactions vary with the number of electrons. However, we believe that the difference between using these two methods becomes unimportant for a higher number of electrons (in the previous analysis [9] very good agreement was found between pairwise interactions as a function of RAM for $N = 7, 8$, even though some discrepancies occurred for $N = 5, 6$).

In Fig. 1A we plot numerical results of exact diagonalization (pluses) and shell model results (circles) for three-quasielectron excitation for $N = 7$. The same is done for $N = 8$ (Fig. 1B); Fig. 2A gives the results for three-quasihole excitations for $N = 7$, and Fig. 2B gives the same results for $N = 8$. We observe very good qualitative agreement between numerical data and the shell model results.

Let us now analyse the spectra of two quasielectrons (quasiholes) and a quasihole (quasielectron). These appear as the second excitation sector for the states containing one quasielectron (quasihole). For $N = 7$ the single quasielectron excitation is in the shell $l = 3.5$, so that in the second excitation sector we have two quasielectrons in the shell $l = 3.5$ and a quasihole in $l = 2.5$. The allowed angular momenta of two quasielectrons are 6, 4, 2, 0, so that the allowed total angular momenta are 0.5, $(1.5)^2$, $(2.5)^3$, $(3.5)^3$, $(4.5)^3$, $(5.5)^2$, $(6.5)^2$, 7.5, 8.5. Again we plot the numerical results together with the results of the shell model (Fig. 3A). First comparison of the numerical results with the non-interacting quasiparticles state suggests that some of the predicted states are missing (or pushed up in energy). In fact, we confirm this in our shell model approach, the main reason for pushing some states up in energy is the large quasielectron-quasihole interaction for $L_{QE-QH} = 1$. It should be noted that other authors [13–15] have interpreted the energy spectra in terms of individual CF excitations. Although no interactions between CF excitations were included, the overlap of the CF state trial wavefunction with the lowest electron Landau level was evaluated. The absence of the $L = 1$ QE-QH state in the first excited sector was attributed to lack of overlap. We simply assume the interaction to be large (we take the numerical result for the lowest energy state with $L = 1$ in the Laughlin state). The same analysis is made for $N = 8$ (Fig. 3B) and for two quasiholes and a quasielectron for $N = 7$ and 8 in Fig. 4A and 4B. In all cases we observe that the qualitative feature of pushing some states up is confirmed by the shell model approach.

There are quantitative discrepancies between the shell model results and the numerical data. Some of them may indicate the presence of another excited states. For example for 2QE-QH excitations we observe that an additional state seems to appear at $l = l_{QE} + 1$, where agreement between our model and the numerical results is the weakest. This is undoubtedly due to our neglect of

the single quasielectron excitation in the next CF Landau level. The “bare” QE energy of this excitation is very close to the “bare” energy of the 2QE-QH state and its angular momentum is $l_{QE} + 1$. Other discrepancies may be a measure of the role of three particle forces which are present in the Chern-Simons approach.

It is interesting to analyse the energy of the ground state of $2/5$ filling. This state can be regarded as one with n_{QE} quasielectrons of the $\nu = 1/3$ state filling the $2l_{QE} + 1$ states of the first excited CF Landau level. The interaction energy of this closed shell configuration of n_{QE} quasielectrons is given by [5]:

$$V(l^{2l+1} L = 0) = \sum_{L'} (2L' + 1) V_{ll}(L'). \quad (5)$$

By adding this energy to $n_{QE}\xi_{QE}$ (sum of single quasielectron energies) we can compare the resulting energy with the exact diagonalization results. The results for $N = 6, 8$ are given in Table I, and they give additional strong support for the “Fermi liquid” shell model picture of CF excitations which we use.

ACKNOWLEDGMENTS

This work was supported in part by Oak Ridge National Laboratory, managed by Lockheed Martin Energy Research Corp. for the US Department of Energy under contract No. DE-AC05-96OR22464. P.S. acknowledges support from NATO High Technology Linkage Grant 930746 and the Foundation for Polish Science. S.N.Y. acknowledges support by the Korea Science and Engineering Foundation. K.S.Y. acknowledges support by the BSRI-95-2412 program of the Ministry of Education, Korea. The authors thank to Drs. G. Canright and A. Kurana for helpful discussions, particular thanks are due to Dr. X. M. Chen for his assistance with the numerical calculations.

-
- [1] F. D. M. Haldane, Phys. Rev. Lett. **51**, 605 (1983).
 - [2] S. He, X. Xie, and F. Zhang, Phys. Rev. Lett. **68**, 3460 (1992).
 - [3] J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989).
 - [4] A. Lopez and E. Fradkin, Phys. Rev. B **44**, 5246 (1991).
 - [5] A. de-Shalit and I. Talmi, *Nuclear Shell Theory*, Academic Press, New York and London 1963.
 - [6] B. I. Halperin, P. A. Lee, and N. Read, Phys. Rev. B **47**, 7312 (1993).
 - [7] G. Fano, F. Ortolani, and E. Colombo, Phys. Rev. B **34**, 2607 (1986).
 - [8] X. M. Chen and J. J. Quinn, Solid State Commun. **92**, 865 (1994).

- [9] S. N. Yi, X. M. Chen, and J. J. Quinn, Phys. Rev. B **53**, xxx (1996).
 [10] M. Johnson and G. Canright, Phys. Rev. B **49**, 2947 (1994).
 [11] See, for example, D. Pines and P. Nozieres, *The Theory of Quantum Liquids*, W. A. Benjamin, Inc, New York 1966.
 [12] M. E. Rose, *Elementary Theory of Angular Momentum*, New York - John Wiley and Sons, London - Chapman and Hall 1957.
 [13] G. Dev and J. K. Jain, Phys. Rev. Lett. **69**, 2843 (1992).
 [14] E. Rezayi and N. Read, Phys. Rev. Lett. **72**, 900 (1994).
 [15] X. E. Wu and J. K. Jain, Phys. Rev. B **51**, 1752 (1995).




FIG. 1. A. Energy spectrum of three quasielectrons for $N = 7$; $l_{QE} = 2.5$ and $L = 1.5, 2.5, 4.5$. V is the interaction energy, i.e. $V = 0$ corresponds to the “bare” energy of three quasielectrons (in all figures energy in units e^2/l_0 where l_0 is the magnetic length for the Laughlin $1/3$ state). B. Same for $N = 8$; $l_{QE} = 3$ and $L = 0, 2, 3, 4, 6$.




FIG. 2. A. Energy spectrum of three quasiholes for $N = 7$; $l_{QH} = 4.5$ and $L = 1.5, 2.5, 3.5, (4.5)^2, 5.5, 6.5, 7.5, 8.5, 10.5$. B. Same for $N = 8$; $l_{QH} = 5$ and $L = 0, 2, 3, 4^2, 5, 6^2, 7, 8, 9, 10, 12$.




FIG. 3. A. Energy spectrum of two quasielectrons and a quasihole for $N = 7$; $l_{QE} = 3.5$, $l_{QH} = 2.5$, and $L = 0.5, (1.5)^2, (2.5)^3, (3.5)^3, (4.5)^3, (5.5)^2, (6.5)^2, 7.5, 8.5$. The significant discrepancy at $L = 4.5$ is undoubtedly associated with the state corresponding to an excited QE in the next CF Landau level which we have neglected. The dashed line in this figure and Fig. 4 indicates the expected position of the “gap”. B. Same for $N = 8$; $l_{QE} = 4$, $l_{QH} = 3$, and $L = 0, 1, 2^3, 3^3, 4^4, 5^3, 6^3, 7^2, 8^2, 9, 10$. The excited single QE state occurs for $L = 5$.




FIG. 4. A. Energy spectrum of two quasiholes and a quasi-electron for $N = 7$; $l_{QE} = 4.5$, $l_{QH} = 3.5$, and $L = 0.5, (1.5)^2, (2.5)^3, (3.5)^3, (4.5)^4, (5.5)^3, (6.5)^3, (7.5)^2, (8.5)^2, 9.5, 10.5$. B. Same for $N = 8$; $l_{QE} = 5$, $l_{QH} = 4$, and $L = 0, 1, 2^3, 3^3, 4^4, 5^4, 6^4, 7^3, 8^3, 9^2, 10^2, 11, 12$.

TABLE I. Comparison of the exact ground state energy of the state of $2/5$ with the shell model results (energy in units e^2/l_0 where l_0 is the magnetic length for the Laughlin $1/3$ state).

	1/3	2/5	
	Exact	Exact	Predicted
$N = 6$	3.871635	4.001568	4.003656
$N = 8$	6.362649	6.521887	6.535434